

Table Adverse events (Carvedilol vs EVL)

Outcome or Subgroup	Studies	Participants	Statistical Method	Effect Estimate
Chest pain requiring medication	1	168	Risk Ratio (M-H, Fixed, 95% CI)	0.03 [0.00, 0.49]
Shortness of breath	3	384	Risk Ratio (M-H, Fixed, 95% CI)	24.57 [4.94, 122.14]
Nausea	2	320	Risk Ratio (M-H, Fixed, 95% CI)	21.52 [2.97, 155.92]
Transient dysphagia	1	168	Risk Ratio (M-H, Fixed, 95% CI)	0.01 [0.00, 0.14]
Orthostatic or symptomatic hypotension	2	320	Risk Ratio (M-H, Fixed, 95% CI)	5.05 [0.60, 42.76]
Impotence	1	152	Risk Ratio (M-H, Fixed, 95% CI)	6.82 [0.36, 129.83]
Dizziness	1	64	Risk Ratio (M-H, Fixed, 95% CI)	2.82 [0.12, 66.82]
Diarrhea	1	64	Risk Ratio (M-H, Fixed, 95% CI)	2.82 [0.12, 66.82]
Hallucinations	1	64	Risk Ratio (M-H, Fixed,	2.82 [0.12, 66.82]

			95% CI)	
Hyperglycaemia	1	64	Risk Ratio (M-H, Fixed, 95% CI)	2.82 [0.12, 66.82]